

Can Stellar Yields Accurately Constrain the Upper Limit to the Initial Mass Function?

Brad K. Gibson

Mount Stromlo & Siding Spring Observatories, Australian National University, Weston Creek P.O., Weston, ACT, Australia 2611

ABSTRACT

Recent determinations of the upper mass limit to the local initial mass function (IMF) claim a value of $m_U = 50 \pm 10 M_\odot$, based upon direct comparisons of the observed oxygen and iron abundances in metal-poor stars with the predicted stellar yields from Type II supernovae (SNe). An unappreciated uncertainty in these analyses is the input physics intrinsic to each SNe grid, and its effect upon stellar nucleosynthesis. We demonstrate how such uncertainties, coupled with the uncertain metal-poor halo star normalization, while allowing us to set a *lower* bound to m_U of $\sim 40 M_\odot$, nullifies any attempt at constraining the *upper* bound.

Subject headings: nuclear reactions, nucleosynthesis, abundances — stars: luminosity, mass function — stars: supernovae — galaxy: evolution

1. Introduction

The initial mass function (IMF), a measure of the distribution of masses, at formation, of a given stellar generation, is one of the key components of galaxy evolution modeling. Despite its importance, accurate determination of the IMF remains one of the most elusive problems in modern astronomy. This elusiveness is manifested both in ongoing attempts to understand its physical underpinnings (e.g. Padoan, Nordlund & Jones 1997), as well as simply characterizing its shape and mass limits from an observational tack (e.g. Kroupa, Tout & Gilmore 1993; Scalo 1986).

In its simplest form, the IMF can be considered a power law of the form $m\phi(m)dm \propto m^{-(1+x)}dm$, where $m\phi(m)dm$ is the number of stars born in the mass interval $m \rightarrow m + dm$. The goal for theorists and observers, alike, is then the determination of the slope of this function x , as well as its upper and lower limits (m_U and m_ℓ , respectively).

While the slope of the IMF, at least in the solar neighborhood (and for masses greater than a few solar masses), would appear to lie somewhere in the range $x \approx 1.3$ (Salpeter 1955) to $x \approx 1.7$ (Kroupa, Tout & Gilmore 1993), and the lower mass limit is close to $m_\ell \approx 0.2 M_\odot$ (Bahcall et al. 1994), the upper mass limit m_U still remains *highly* uncertain. Even a cursory examination of the literature corroborates this point, with values in the range $m_U \approx 20 \rightarrow 200 M_\odot$ suggested by a variety of direct and indirect techniques (e.g. Maeder & Meynet 1989; Klapp & Corona-Galinda 1990; Pagel et al. 1992; Maeder 1992; Massey et al. 1995; Kudritzki 1997).

A hybrid approach to determining m_U , combining predictions of the *theoretical* yields from Type II supernovae (SNe), with the *observed* abundances in the old metal-poor stars (i.e. those which bear the clear imprint of yield “pollution” from these same SNe, with no “dilution” from Type Ia SNe, whose progenitor lifetimes are considerably longer than the Type II timescales), has been the subject a recent series of papers (Tsujiimoto et al. 1995, 1997; Yoshii, Tsujiimoto & Nomoto 1996). The premise here is that because Type II SNe α -element (e.g. O, Mg, Ne) yields are a strong function of progenitor mass, whereas products of explosive burning (e.g. Fe, Si, Ca) are less so, the IMF-weighted average of their ratios must necessarily also depend strongly upon x and m_U . Tying these yield “averages” to the halo abundances then, in principle, provides a unique indirect probe on the

upper mass limit to the IMF.¹

Following this technique, Tsujiimoto et al. (1997) recently concluded that the upper mass limit to the IMF in the solar neighborhood is $m_U = 50 \pm 10 M_\odot$. To do so, they made explicit use of the Tsujiimoto et al. (1995) compilation of Type II SNe yields. What was not fully appreciated in their study though was just how dependent their result was to this *particular* yield compilation and the adopted halo abundance normalization. It is to this lack of appreciation that our current study is addressed.

After providing a minimal introduction to the model ingredients in Section 2.1, we demonstrate in Sections 2.2 and 2.3 that this technique results in $m_U = 50 \pm 10 M_\odot$ *only* for the Tsujiimoto et al. (1995) yields, combined with a halo normalization of $[O/Fe]_h = +0.41$. Duplicating the analysis with “competing” yield compilations which sample a wide variety of convection and mass-loss treatments (e.g. Woosley & Weaver 1995; Langer & Henkel 1995; Arnett 1996), clearly demonstrates that Tsujiimoto et al. (1997) have significantly underestimated the uncertainty associated with their determination of m_U . Our results are summarized in Section 3.

2. Analysis

2.1. The Basic Formalism

The observed oxygen-to-iron abundance ratio in halo dwarfs and giants, $[O/Fe]_h$, can be linked (see Tsujiimoto et al. 1997 – hereafter, T97 – for details) to the theoretical Type II SNe yields m_O^{ej} and m_{Fe}^{ej} via

$$\left(\frac{O}{Fe}\right)_h = \left(\frac{O}{Fe}\right)_\odot \times 10^{[O/Fe]_h} = \frac{\int_{10}^{m_U} m_O^{ej} m^{-(1+x)} dm}{\int_{10}^{m_U} m_{Fe}^{ej} m^{-(1+x)} dm}, \quad (1)$$

where the coefficient $(O/Fe)_\odot \equiv 7.55$ is the solar (meteoritic) mass fraction ratio from Anders & Grevesse (1989), and x is the slope of the IMF (recall Section 1).

By specifying the low metallicity “plateau” value for $[O/Fe]_h$ (e.g. Figure 11 of Timmes, Woosley & Weaver 1995; Figure 2 of Bessell, Sutherland & Ruan

¹Tsujiimoto et al. (1997) extend this “hybrid” approach to simultaneously constrain the IMF mass limits m_U and m_ℓ , as well as the slope x . We shall only be concerned with the m_U determination in what follows, primarily for brevity, but also because the lower mass constraint rest squarely upon uncertain photometric calibrations.

1991), and the IMF slope x , equation 1 provides a unique upper mass limit m_U , for a given *Type II SNe yield compilation*.

2.1.1. Ingredients

Essentially, there are only two input ingredients which interest us, in what follows. First, the source of Type II SNe yields (right-hand side of equation 1), and second, the halo’s “plateau” $[O/Fe]_h$ (left-hand side of equation 1). Let us briefly comment on the latter ingredient first.

There is no debate as to the reality of the α -element overabundance seen in halo dwarfs and giants (i.e. $[O/Fe]_h \gtrsim +0.2$), but there does remain a factor of ~ 2 uncertainty as to its asymptotic “plateau” value for $[Fe/H] \lesssim -3$. T97 adopt $[O/Fe]_h = +0.41^2$ but it is apparent that this depends somewhat upon the sample selection. Visual inspection of Figure 11 of Timmes et al. (1995), Figure 2 of Bessell et al. (1991), and Figure 1 of T97, each drawn from slightly different sources, shows that the plateau lies roughly between $[O/Fe] \approx +0.4 \rightarrow +0.6$; i.e. T97’s value of $+0.41$ appears to lie on the lower end of the plateau “distribution”. T97 did not demonstrate how the predicted m_U changes as a function of this halo “normalization”, a point to which we return in Section 2.3, where we duplicate their analysis using $[O/Fe]_h = +0.60$ (i.e. that favored by Bessell et al. 1991).

The Type II SNe yields are the primary input into equation 1. T97 adopt their earlier 1995 yield compilation (T95, hereafter), which itself is an offshoot of the Thielemann, Nomoto & Hasimoto (1996) models. The solid curves in Figure 1 represent the T95 oxygen and iron yields adopted in the T97 analysis.

It is readily apparent though, as even a cursory

²T97’s claim to adopt $[O/Fe]_h = +0.41$ is not entirely correct, as their code does not follow O and Fe, per se, but only the isotopes ^{16}O and ^{56}Fe , thereby underestimating the true iron yield by $\sim 10\%$ (and oxygen by a smaller amount). This has the effect that instead of using $(O/Fe)_\odot = 7.55$ in equation 1, T97 use $(O/Fe)_\odot = 8.20$ (i.e. $(^{16}O/^{56}Fe)_\odot = 8.20$). By itself, this is not a problem; the problem arises in that T97 then normalize equation 1 to the observed halo ratio $[O/Fe]_h = +0.41$ - this observed ratio, though, reflects the *total* O and Fe abundances, and *not* just the isotopes ^{16}O and ^{56}Fe . In practice, what this means is that T97 use $(O/Fe)_h = 21.1$ (recall equation 1), which corresponds to $[O/Fe]_h = +0.45$, and *not* $+0.41$. To truly normalize their models to $[O/Fe]_h = +0.41$, T97 should have used $(O/Fe)_h = 19.4$ and included the total O and Fe for the yields in the right-hand side of equation 1.




Fig. 1.— Oxygen (upper four curves) and iron (lower three curves) yields as a function of progenitor mass, for the primary yield compilations considered here – T95=Tsujimoto et al. (1995); LH95=Langer & Henkel (1995); A96=Arnett (1996); WW95=Woosley & Weaver (1995); M92=Maeder (1992). The models of M92, LH95 and A96 are only evolved up to the completion of oxygen burning, hence an iron yield is not directly associated with either grid. We indirectly associate Arnett’s (1991) iron yields with his newer 1996 oxygen yields. See text for details.

glance at Figure 1 should show, that there still remain *substantial* uncertainties in the yield computations for Type II SNe. These differences have already been described in detail by Arnett (1995), Langer (1997), and Gibson, Loewenstein & Mushotzky (1997), to name just a few.

T97 assume that these differences can be reconciled by considering what they (mistakenly) assume is the maximal deviation from their canonical (T95) yields. This deviation is taken to be a mass-independent 30% excess added to their oxygen yields; iron, though, is assumed to be invariant. T97 justify this 30% excess as the “extreme”, since Woosley & Weaver’s (1995) (WW95, hereafter) $[Fe/H]=+0.0$ oxygen yields are roughly 30% greater than those of

T95 (also $[\text{Fe}/\text{H}]=+0.0$).³ Herein lies several (intertwined) problems.

First, solar metallicity yields are inappropriate for the discussion at hand; T97 attempt to constrain m_{U} by comparing the abundance ratios in metal-poor stars (i.e. $[\text{Fe}/\text{H}] \lesssim -2$) with the yields predicted from $[\text{Fe}/\text{H}]=+0.0$ Type II SNe models. A more appropriate course of action is to consider a much lower metallicity grid of models. Because T95 do not compute $[\text{Fe}/\text{H}] < +0.0$ models, T97 could not explicitly consider metallicity effects.

If we compare (properly) T95’s oxygen yields with WW95’s sub-solar metallicity grid, the 30% “excess” adopted by T97 is no longer relevant. Figure 1 shows that WW95’s $[\text{Fe}/\text{H}]=-4.0$ models have oxygen yields virtually indistinguishable to the $[\text{Fe}/\text{H}]=+0.0$ models of T95, except for $m \gtrsim 35 M_{\odot}$, where WW95 now lies systematically *below* T95.

These subtle differences in the WW95 and T95 oxygen yields are interesting in their own right (e.g. Langer 1997), and are certainly encapsulated by the 30% error budget adopted by T97. Unfortunately, a more fundamental flaw made by T97 was in assuming that *all* of the uncertainty in the oxygen yields could be accounted for by this 30% factor. The problem with assuming that the T95 oxygen yields form a lower envelope, with WW95 forming the upper, can best be appreciated by referring to the two other primary sources of SNe models, besides those of T95 and WW95 – i.e. Langer & Henkel (1995, hereafter LH95) and Arnett (1996, hereafter A96).

In Figure 1, we can see that both LH95 and A96 agree (roughly) with one another,⁴ *but* lie a factor of ~ 3 below the T95 and WW95 predictions. This factor of three uncertainty in the oxygen yields is a far cry from the 30% uncertainty advocated by T97, but is entirely in keeping with that found by Langer (1997). This latter reference is an invaluable resource for those interested in properly assessing the origin of the large uncertainties in different modelers’ oxygen yields. In the same vein, a useful comparison of the primary differences in the model input physics

can be found in Table 2 of Gibson, Loewenstein & Mushotzky (1997). At this point, we are *not* advocating one grid of oxygen yields over another, simply drawing attention to the fact that T97 have underestimated its uncertainty.

Figure 1 reflects the present-day state-of-the-art, as far as oxygen yields goes, and can be considered to be the modern equivalent of Wang & Silk’s (1993) Figure 1. The similarity of the Arnett (1978), Woosley & Weaver (1986), and Thielemann, Nomoto & Hashimoto (1994) oxygen yields, reflected in the latter figure, is not in dispute, but it should also be readily apparent (from our Figure 1) that supplementing the current analogs of these older models (i.e. A96, WW95, and T95, respectively) with the new models of LH95 demonstrates that the agreement is no better than a factor of $\sim 2 \rightarrow 3$ at a given initial mass, in agreement with that found by Langer (1997).

Figure 2 parallels that of Figure 1, but restricts itself to the oxygen yields from the published compilations of Arnett (1978,1991,1996), hereafter A78, A91, and A96, respectively. The subscript ‘He’ or ‘ZAMS’ denotes the initial evolutionary state of the stellar models – pure Helium core or zero age main sequence. The former implies that a ZAMS mass-He core mass relation has been applied *a posteriori*, similar to the models of T95, whereas the latter avoid such complications by modeling the evolution self-consistently. We include a parallel analysis using each of the Arnett generations as (i) A78 has already played a large part in the analysis of Wang & Silk (1991), and for completeness sake should therefore be included here; (ii) A78 and A91 are identical in all respects, save the adopted He core mass-ZAMS mass relation, a useful comparison in and of itself; (iii) A96_{He} and A96_{ZAMS} are similar in all respects, save initial evolutionary state, thereby allowing us to better appreciate this crucial difference in the modeling. Langer (1997) has previously alluded to the danger of comparing yields from models evolved from the ZAMS versus those evolved from He cores; we are now in a position to quantify this “danger”, as applied to the determination of the IMF upper mass limit.

A further assumption inherent to the T97 analysis is that the iron yields are invariant and best represented by those of T95. Iron predictions are particularly problematic as the exact location of the mass-cut leads to enormous uncertainties (Thielemann et al. 1996). For the sake of self-consistency in the input physics though, we have not made the same *a priori*

³Actually, WW95’s $[\text{Fe}/\text{H}]=+0.0$ oxygen yields (their favored Case B energetics) are greater than those of T95 *only* for $m \lesssim 30 M_{\odot}$.

⁴The treatment of convection, the adopted $^{12}\text{C}(\alpha,\gamma)^{16}\text{O}$ reaction rate, and the initial evolutionary state of the models, were all significantly different in T95, than in either LH95 or A96. LH95 also considered sub-solar metallicity models and self-consistent mass-loss, both of which the others did not.




Fig. 2.— Oxygen yields as a function of progenitor mass, from the four compilations of Arnett (1978, 1991, 1996). Only the A96_{ZAMS} grid was evolved self-consistently from the zero age main sequence (ZAMS), while the remainder were evolved helium cores with an assumed *a posteriori* ZAMS mass-He core mass relation.

assumption regarding iron’s invariance, and have attempted to match the oxygen predictions with iron from the same grid. This is easy in the case of T95 and WW95, as they follow the evolution right through core-collapse and explosive nucleosynthesis. Of the different Arnett grids described herein, only A91 included a specific entry for iron, so that is used for each of A78, A91, and A96, hereafter. Proper self-consistency with Arnett’s yields will be restored after he applies core collapse and explosive nucleosynthesis to the new A96 model grid.

We are now in a position to anticipate some of the conclusions of Section 2.2. To do so, we weight the mass-dependent oxygen and iron yields of Figures 1 and 2 by some canonical IMF, which we will take to be of the slope favored by Salpeter (1955) - i.e. $x = 1.35$ - with Type II SNe progenitors assumed to span the mass range $10 \rightarrow 50 M_{\odot}$. The resulting mean IMF-weighted masses of oxygen and iron (and their logarithmic ratios relative to the solar ratio) - i.e. $\langle O \rangle$, $\langle Fe \rangle$, and $\langle [O/Fe] \rangle$ - for each of the

yield “pairs” discussed thus far, are listed in Table 1. The subscripts ZAMS and He refer to the initial evolutionary state of the models - i.e. zero age main sequence or simple helium cores.

The factor of ~ 3 uncertainty due to oxygen (and to a lesser extent, that due to iron) is immediately apparent in columns 3 and 4, their combination leading to a factor of ~ 2 uncertainty in the IMF-weighted $[O/Fe]$.

Recalling that T97, using the T95 yields, assumed a metal-poor normalization for equation 1 of $[O/Fe]_h \approx +0.4$, and subsequently found $m_U \approx 50 M_{\odot}$, it should not be surprising to see that our entry in Table 1 for T95, generated using $m_U = 50 M_{\odot}$, has an IMF-weighted $[O/Fe]$ similar to T97’s adopted halo value. Conversely, the A96 entries lie substantially below this halo value. Because $[O/Fe]$ increases with increasing progenitor mass (see Figure 1), it should be apparent that to recover an $[O/Fe] \approx +0.4$, with the same IMF slope, will require a greater m_U .

2.2. Quantifying m_U ’s Dependence Upon Yield Source

For a given halo “plateau” normalization $[O/Fe]_h$, the various yield compilations of Figures 1 and 2 (and Section 2.1.1) can now be used in conjunction with equation 1 to determine how the upper mass limit m_U varies as a function of IMF slope x . The results, for a variety of models, are shown in Figure 3. Let us restrict ourselves, for the time being, to the six curves which were generated assuming the default halo normalization of $[O/Fe] = +0.41$.

The first model to note from Figure 3 is that given by the solid curve, which effectively recovers the results of T97⁵ - i.e. $m_U \approx 48 \rightarrow 62 M_{\odot}$, for reasonable IMF power-law slopes spanning $x \approx 1.3$ (Salpeter 1955) to $x \approx 1.7$ (Kroupa et al. 1993). Again, recalling our comments at the end of Section 2.1.1, this low predicted value for m_U was already anticipated, based upon the T95 yields possessing the greatest IMF-weighted $\langle [O/Fe] \rangle$, for the “standard” IMF employed in constructing Table 1.

The most important result to take from Figure 3, though, is not so much the behavior of the T95 curve *alone*, but the parallel behavior for the models employing the alternate Type II yields - i.e. those de-

⁵The footnote to Section 2.1.1 clarifies why there remains a residual $\sim 10\%$ offset between the two.

TABLE 1
IMF-WEIGHTED YIELDS^a

Source ^b	[Fe/H] ^c	< O > ^d	< Fe > ^d	< [O/Fe] >
T95 _{He}	+0.0	1.810	0.091	+0.42
M92 _{ZAMS}	-1.3	1.508	n/a	n/a
WW95 _{ZAMS}	-4.0	1.445	0.087	+0.34
LH95 _{ZAMS}	-1.0	0.841	n/a	n/a
A78 _{He} ^O +A91 _{He} ^{Fe}	+0.0	1.711	0.068 ^e	+0.52
A91 _{He} ^O +A91 _{He} ^{Fe}	+0.0	1.189	0.068 ^e	+0.37
A96 _{He} ^O +A91 _{He} ^{Fe}	+0.0	0.983	0.068 ^e	+0.28
A96 _{ZAMS} ^O +A91 _{He} ^{Fe}	+0.0	0.678	0.068 ^e	+0.12

^a IMF slope $x = 1.35$, over the range $10 \rightarrow 50 M_{\odot}$.

^b Source of oxygen yields.

^c Metallicity of models from which oxygen yields were derived.

^d IMF-weighted yield mass, in $M_{\odot}$.

^e Arnett (1991) [Fe/H]=+0.0 iron yields.

noted WW95 and A96,⁶ in the legend to the figure. For a Salpeter (1955) IMF slope (i.e. $x = 1.35$), the predicted upper mass limit for the T95 yields was $m_U \approx 49 M_{\odot}$. For comparison, the corresponding m_U for the WW95, A78 and A96_{ZAMS} yields is 61, 35, and $131 M_{\odot}$, respectively, with the A91 and A96_{He} values lying between A78 and A96_{ZAMS}. Because the dependence of m_U upon IMF slope x is stronger in the non-T95 yields (especially for A96_{ZAMS}), for the Kroupa et al. (1993) slope (i.e. $x = 1.7$) m_U is more strongly affected. The value favored by the T95 compilation (i.e. $62 M_{\odot}$) is replaced, in this case, by values ranging from $\sim 95 M_{\odot}$, for WW95,⁷ to $\gtrsim 200 M_{\odot}$, for A96_{ZAMS}.

⁶We have not shown the results for the LH95 oxygen yields in Figure 3, as they parallel closely those of A96_{ZAMS}.

⁷Because the maximum mass considered by WW95 was $40 M_{\odot}$, we make the (perhaps mistaken) assumption that linear extrapolation beyond the grid extreme holds. For example, such an assumption leads to a predicted oxygen yield of $14 M_{\odot}$ for an $m = 100 M_{\odot}$ star. If we were to arbitrarily double this yield to $\sim 30 M_{\odot}$, and set the iron yield to a constant $0.075 M_{\odot}$ for $m \geq 35 M_{\odot}$, we would find an approximate 5%/15% reduction in the predicted m_U , for IMF slopes $x = 1.3/1.6$. Even more extreme, if we increase the same oxygen yield to $50 M_{\odot}$, and set the iron yield to zero for $m \geq 35 M_{\odot}$, we would find an approximate 15%/30% reduction in the predicted m_U , for IMF slopes $x = 1.3/1.6$. Obviously, self-consistent yield predictions for the WW95 grid, for masses $m > 40 M_{\odot}$, are required, although the previous examples do provide a feel for maximum effect involved.

To summarize, *only if one assumes that the T95 yields are the definitive representation of low-metallicity massive star nucleosynthetic yields can one conclude that the upper mass limit to the solar neighborhood IMF is $m_U = 40 \rightarrow 60 M_{\odot}$* . By adopting, in turn, the three other primary yield sources (i.e. WW95, LH95, and A96), we sample a much fairer representative input physics “parameter space”, leading to the more conservative result that we are able to constrain the *lower* bound to the upper mass limit to be $\sim 40 M_{\odot}$, but that the *upper* bound remains effectively unconstrained. If we force the Salpeter (1955) IMF slope (i.e. $x = 1.35$) to hold, the uncertainty in the yields still only allows us to constrain m_U to the range $\sim 40 \rightarrow 140 M_{\odot}$. *This yield-dependence of m_U ’s determination was not appreciated in the original analysis by T97.*

It is interesting to question how one might recover a result of $m_U \approx 40 \rightarrow 60 M_{\odot}$, when adopting the A96_{ZAMS} yields. Assuming the oxygen as inviolate, it is apparent from inspection of Table 1 that we are required to reduce the IMF-weighted iron yield by a factor of ~ 2 . This cannot be of the form of a blanket, mass-independent reduction, since we are constrained by the observations of SNe 1987a and 1993j (Thielemann et al. 1996). Only by setting the iron yield to zero for $m \leq 12 M_{\odot}$ and $m \geq 30 M_{\odot}$ (i.e. retaining the 15, 20, and $25 M_{\odot}$ iron pre-




Fig. 3.— Upper mass limit m_U , as a function of IMF slope x , for the Type II SNe yield compilations of Figures 1 and 2.

dictions of A91, but ignoring the contribution from other masses), can we reduce the A96_{ZAMS} prediction for m_U from $\sim 130 \rightarrow 200 M_\odot$ to that found using T95 (i.e. $m_U \approx 40 \rightarrow 60 M_\odot$).

In passing, we note that linear interpolation for both oxygen and iron were assumed in equation 1. If one adopts a logarithmic interpolation, as Yoshii et al. (1996) did, the predicted values of m_U would be reduced by $\sim 10 \rightarrow 25\%$, from those shown in Figure 3.

2.3. Quantifying m_U 's Dependence Upon the Halo Normalization

T97 did not consider how their conclusions might be influenced by the adopted mean “halo” $[O/Fe]_h$ (equation 1). Recall that the solid curve in Figure 3 supported $m_U \approx 48 \rightarrow 62 M_\odot$, for IMF slopes $x \approx 1.3 \rightarrow 1.7$, for the T95 yields and a normalization $[O/Fe]_h = +0.41$.

If instead of adopting T97's favored halo normalization, we took that from Bessell et al. (1991) - i.e. $[O/Fe]_h = +0.60$ - we would recover the “T95;+0.60” curve of Figure 3. In other words, this ~ 0.2 dex (i.e. $\sim 60\%$) higher normalization, for the T95 yields, increases the predicted m_U by $\sim 60\%$ (from $48 M_\odot$ to

$75 M_\odot$, for $x = 1.3$) to $\sim 140\%$ (from $62 M_\odot$ to $148 M_\odot$, for $x = 1.7$). The same conclusion is reached when we adopt the WW95 yields, as shown by the “WW95;+0.60” curve of Figure 3. A general rule-of-thumb for Salpeter (1955) IMF slopes (i.e. $x = 1.35$) is that a given percentage increase in the halo normalization, is accompanied by the same percentage increase in the predicted m_U , regardless of yield source. For slopes $x \approx 1.7$, the increase in m_U with increasing normalization is generally $\sim 2 \rightarrow 3\times$ greater.

If we simply take the halo normalization to be $[O/Fe]_h = +0.5 \pm 0.1$, and accept T97's allowable range of IMF slopes (i.e. $x \approx 1.3 \rightarrow 1.6$), Figure 3 would lead us to conclude that $m_U \approx 40 \rightarrow 140 M_\odot$ better represents the valid range of m_U , for the T95 yields— we feel that this would have been a more realistic range than the $m_U \approx 40 \rightarrow 60 M_\odot$ claimed by T97.

3. Summary

T97 have recently revitalized interest in using IMF-weighted Type II SNe yields as a direct probe of said IMF's upper mass limit m_U , by comparison with the observed abundance ratios in metal-poor Galactic stars. The beauty of this technique lies partially in its simplicity – for a given IMF slope, there is effectively only one free parameter – the yield source. Adopting the T95 yields, T97 found $m_U = 40 \rightarrow 60 M_\odot$, for reasonable IMF slopes (i.e. $x = 1.3 \rightarrow 1.6$).

The primary concern we have regarding T97's analysis lies in their underlying assumption that *all* of the uncertainty in the stellar model physics could be encapsulated in a 30% (0%) error budget for oxygen (iron). It should be obvious from Figures 1 and 2, and Langer (1997), for example, that this assumption is incorrect, and that a more realistic error budget would allow for up to an order of magnitude greater leeway. Stellar models are simply not developed to the level that is inherently assumed by T97 – convection, overshooting, mass-loss, reaction rates, metallicity, C/O-core masses, fallback onto the remnant – each conspire to increase the uncertainties to the degree reflected by the yields shown in Figures 1 and 2.

A secondary concern is T97's inherent assumption that the halo normalization $[O/Fe]_h = +0.41$ has no associated uncertainty. Since values as high as $[O/Fe] = +0.6$ are still favored by some, this $\sim 60\%$ uncertainty should be taken into account. For a

Salpeter (1955) slope, there is (roughly) a one-to-one correspondence between the halo normalization uncertainty and the corresponding predicted upper mass limit uncertainty.

While we *do* agree with T97 that the *lower* limit to m_U is $\sim 40 M_\odot$ (or $\sim 30 M_\odot$, if we adopt the extreme A78 yields), our more realistic exploration of input physics “space” demonstrates that we simply cannot constrain the *upper* limit to any useful accuracy. Taken together, we can only conclude that, by this technique alone, $m_U \gtrsim 40 M_\odot$, for IMF slopes $x = 1.3 \rightarrow 1.6$. Fixing the IMF slope to that of Salpeter (1955), we can only constrain m_U to lie somewhere between $\sim 40 M_\odot$ and $\sim 140 M_\odot$. Finally, it would appear to be difficult to reconcile any $m_U \lesssim 100 M_\odot$ with either the A96ZAMS or WW95;+0.60 halo normalization grids.

While promising (provided existing discrepancies in Type II SNe yields are eliminated), at the present time, unfortunately, this technique, by itself, does not substantially improve or constrain our understanding of the upper mass limit to the solar neighborhood IMF.

We wish to thank Takuji Tsujimoto and Sylvia Becker for a number of helpful correspondences. BKG acknowledges the financial support of an NSERC Postdoctoral Fellowship.

REFERENCES

- Arnett, D. 1978, ApJ, 219, 1008 (A78)
- Arnett, D. 1991, Frontiers of Stellar Evolution, ed. D.L. Lambert, ASP Conf. Series, 389 (A91)
- Arnett, D. 1995, ARA&A, 33, 115
- Arnett, D. 1996, Supernovae and Nucleosynthesis, Princeton: Princeton Univ. Press (A96)
- Bahcall, J.N., Flynn, C., Gould, A. & Kirhakos, S. 1994, ApJ, 435, L51
- Bessell, M.S., Sutherland, R.S. & Ruan, K. 1991, ApJ, 383, L71
- Gibson, B.K., Loewenstein, M. & Mushotzky, R.F. 1997, MNRAS, 290, 623
- Klapp, J. & Corona-Galindo, M.G. 1990, Rev. Mex. Astron. Astrofis., 21, 317
- Kroupa, P., Tout, C.A. & Gilmore, G. 1993, MNRAS, 262, 545
- Kudritzki, R.P. 1997, Stellar Astrophysics for the Local Group: A First Step to the Universe (VIII Canary Islands Winter School), in press
- Langer, N. 1997, The History of the Milky Way and its Satellite System, eds. A. Burkert, D.H. Hartmann & S.R. Majewski, ASP Conf. Series, 169
- Langer, N. & Henkel, C. 1995, Space Sci. Rev., 74, 343 (LH95)
- Maeder, A. 1992, A&A, 264, 105 (Erratum: 1993, A&A, 268, 833)
- Maeder, A. & Meynet, G. 1989, A&A, 210, 155
- Massey, P., Johnson, K.E. & Degioia-Eastwood, K. 1995, ApJ, 454, 151
- Padoan, P., Nordlund, Å.P. & Jones, B.J.T. 1997, MNRAS, 288, 145
- Pagel, B.E.J., Simonson, E., Terlevich, R. & Edmunds, M. 1992, MNRAS, 255, 325
- Salpeter, E.E. 1955, ApJ, 121, 161
- Scalo, J.M. 1986, Fund. Cosmic Phys., 11, 1
- Thielemann, F.-K., Nomoto, K. & Hashimoto, M. 1994, Supernovae, eds. J. Audouze, S.R. Bludman, R. Mochkovitch & J. Zinn-Justin, Amsterdam: Elsevier, 629
- Thielemann, F.-K., Nomoto, K. & Hashimoto, M. 1996, ApJ, 460, 408
- Timmes, F.X., Woosley, S.E. & Weaver, T.A. 1995, ApJS, 98, 617
- Tsujimoto, T., Nomoto, K., Yoshii, Y., Hashimoto, M., Yanagida, S. & Thielemann, F.-K. 1995, MNRAS, 277, 945 (T95)
- Tsujimoto, T., Yoshii, Y., Nomoto, K., Matteucci, F., Thielemann, F.-K. & Hashimoto, M. 1997, ApJ, 483, 228 (T97)
- Wang, B. & Silk, J. 1993, ApJ, 406, 580
- Woosley, S.E. & Weaver, T.A. 1986, Radiation Hydrodynamics, eds. D. Mihalas & K.H. Winkler, Dordrecht: Reidel, 91

Woosley, S.E. & Weaver, T.A. 1995, ApJS, 101, 181
(WW95)

Yohii, Y., Tsujimoto, T. & Nomoto, K. 1996, ApJ,
462, 266